


Cervical Cancer Screening and Management Recommendations for Individuals >65 Years of Age^a With a Cervix

[Click here for topics associated with this algorithm](#)



^aScreening is not recommended for individuals who are posthysterectomy with no history of cervical cancer or CIN2+ in the past 25 years.

^bAdequate screening is defined as 2 consecutive negative primary HPV tests, 2 consecutive negative cotests, or 3 consecutive cytology tests within the past 10 years. Most recent test must be within the recommended interval (3-5 years) for the test used.

^cCIN2+ is defined as cervical intraepithelial neoplasia grade 2 or more severe disease.

References

Fontham ETH, Wolf AMD, Church TR, et al. Cervical cancer screening for individuals at average risk: 2020 guideline update from the American Cancer Society. *CA Cancer J Clin.* 2020;70(5):321-346.

Huh WK, Ault KA, Chelmow D, Davey DD, et al. Use of primary high-risk human papillomavirus testing for cervical cancer screening: interim clinical guidance. *Gynecol Oncol.* 2015;136(2):178-182.

Massad S, Einstein MH, Huh WK, et al. 2012 ASCCP consensus guidelines conference. 2012 updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors *Obstet Gynecol.* 2013;121(4):829-846.

The American Society for Colposcopy and Cervical Pathology. [Cervical cancer screening recommendations, 2014](#). [Updated: Aug 2014; Accessed: Apr 2020]